

**Table 2. 2002 Criteria for Potentially Inappropriate Medication Use in Older Adults: Independent of Diagnoses or Conditions**

DRUG	CONCERN	SEVERITY RATING (HIGH OR LOW)
Propoxyphene (Darvon) and combination products (Darvon with ASA, Darvon-N, and Darvocet-N)	Offers few analgesic advantages over acetaminophen, yet has the adverse effects of other narcotic drugs.	Low
Indomethacin (Indocin and Indocin SR)	Of all available nonsteroidal anti-inflammatory drugs, this drug produces the most CNS adverse effects.	High
Pentazocine (Talwin)	Narcotic analgesic that causes more CNS adverse effects, including confusion and hallucinations, more commonly than other narcotic drugs. Additionally, it is a mixed agonist and antagonist.	High
Trimethobenzamide (Tigan)	One of the least effective antiemetic drugs, yet it can cause extra-pyramidal adverse effects.	High
Muscle relaxants and antispasmodics: methocarbamol (Robaxin), carisoprodol (Soma), chlorzoxazone (Paraflex), metaxalone (Skelaxin), cyclobenzaprine (Flexeril), and oxybutynin (Ditropan). Do not consider the extended-release Ditropan XL.	Most muscle relaxants and antispasmodic drugs are poorly tolerated by elderly patients, since these cause anticholinergic adverse effects, sedation, and weakness. Additionally, their effectiveness at doses tolerated by elderly patients is questionable.	High
Flurazepam (Dalmane)	This benzodiazepine hypnotic has an extremely long half-life in elderly patients (often days), producing prolonged sedation and increasing the incidence of falls and fracture. Medium- or short-acting benzodiazepines are preferable.	High
Amitriptyline (Elavil), chlordiazepoxide-amitriptyline (Limbital), and perphenazine-amitriptyline (Triavil)	Because of its strong anticholinergic and sedation properties, amitriptyline is rarely the antidepressant of choice for elderly patients.	High
Doxepin (Sinequan)	Because of its strong anticholinergic and sedating properties, doxepin is rarely the antidepressant of choice for elderly patients.	High
Meprobamate (Miltown and Equanil)	This is a highly addictive and sedating anxiolytic. Those using meprobamate for prolonged periods may become addicted and may need to be withdrawn slowly.	High
Doses of short-acting benzodiazepines: doses greater than lorazepam (Ativan), 3 mg; oxazepam (Serax), 60 mg; alprazolam (Xanax), 2 mg; temazepam (Restoril), 15 mg; and triazolam (Halcion), 0.25 mg	Because of increased sensitivity to benzodiazepines in elderly patients smaller doses may be effective as well as safer. Total daily doses should rarely exceed the suggested maximums.	High
Long-acting benzodiazepines: chlordiazepoxide (Librium), chlordiazepoxide-amitriptyline (Limbital), clidinium-chlordiazepoxide (Librax), diazepam (Valium), quazepam (Doral), halazepam (Paxipam), and chlorazepate (Tranzone)	These drugs have a long half-life in elderly patients (often several days), producing prolonged sedation and increasing the risk of falls and fractures. Short- and intermediate-acting benzodiazepines are preferred if a benzodiazepine is required.	High
Disopyramide (Norpace and Norpace CR)	Of all antiarrhythmic drugs, this is the most potent negative inotrope and therefore may induce heart failure in elderly patients. It is also strongly anticholinergic. Other antiarrhythmic drugs should be used.	High
Digoxin (Lanoxin) (should not exceed >0.125 mg/d except when treating atrial arrhythmias)	Decreased renal clearance may lead to increased risk of toxic effects.	Low
Short-acting dipyridamole (Persantine). Do not consider the long-acting dipyridamole (which has better properties than the short-acting in older adults) except with patients with artificial heart valves.	May cause orthostatic hypotension.	Low
Methyldopa (Aldomet) and methyldopa-hydrochlorothiazide (Aldoril)	May cause bradycardia and exacerbate depression in elderly patients.	High
Reserpine at doses >0.25 mg	May induce depression, impotence, sedation, and orthostatic hypotension.	Low
Chlorpropamide (Diabinese)	It has a prolonged half-life in elderly patients and could cause prolonged hypoglycemia. Additionally, it is the only oral hypoglycemic agent that causes SIADH.	High
Gastrointestinal antispasmodic drugs: dicyclomine (Bentyl), hyoscyamine (Levsin and Levsinex), propantheline (Pro-Banthine), belladonna alkaloids (Donnatal and others), and clidinium-chlordiazepoxide (Librax)	GI antispasmodic drugs are highly anticholinergic and have uncertain effectiveness. These drugs should be avoided, especially for long-term use.	High

(continued)

**Table 2. 2002 Criteria for Potentially Inappropriate Medication Use in Older Adults: Independent of Diagnoses or Conditions (continued)**

DRUG	CONCERN	SEVERITY RATING (HIGH OR LOW)
Anticholinergics and antihistamines: chlorpheniramine (Chlor-Trimeton), diphenhydramine (Benadryl), hydroxyzine (Vistaril and Atarax), cyproheptadine (Perlactin), promethazine (Phenergan), tripeleonnamine, dexchlorpheniramine (Polaramine)	All nonprescription and many prescription antihistamines may have potent anticholinergic properties. Nonanticholinergic antihistamines are preferred in elderly patients when treating allergic reactions.	High
Diphenhydramine (Benadryl)	May cause confusion and sedation. Should not be used as a hypnotic, and when used to treat emergency allergic reactions, it should be used in the smallest possible dose.	High
Ergot mesyloids (Hydergine) and cycloandelate (Cyclospasmol)	Have not been shown to be effective in the doses studied.	Low
Ferrous sulfate >325 mg/d	Doses >325 mg/d do not dramatically increase the amount absorbed but greatly increase the incidence of constipation.	Low
All barbiturates (except phenobarbital) except when used to control seizures	Are highly addictive and cause more adverse effects than most sedative or hypnotic drugs in elderly patients.	High
Meperidine (Demerol)	Not an effective oral analgesic in doses commonly used. May cause confusion and has many disadvantages to other narcotic drugs.	High
Ticlopidine (Ticlid)	Has been shown to be no better than aspirin in preventing clotting and may be considerably more toxic. Safer, more effective alternatives exist.	High
Ketorolac (Toradol)	Immediate and long-term use should be avoided in older persons, since a significant number have asymptomatic GI pathologic conditions.	High
Amphetamines and anorexic agents	These drugs have potential for causing dependence, hypertension, angina, and myocardial infarction.	High
Long-term use of full-dosage, longer half-life, non-COX-selective NSAIDs: naproxen (Naprosyn, Avapro, and Aleve), oxaprozin (Daypro), and piroxicam (Feldene)	Have the potential to produce GI bleeding, renal failure, high blood pressure, and heart failure.	High
Daily fluoxetine (Prozac)	Long half-life of drug and risk of producing excessive CNS stimulation, sleep disturbances, and increasing agitation. Safer alternatives exist.	High
Long-term use of stimulant laxatives: bisacodyl (Dulcolax), cascara sagrada, and Neoloid except in the presence of opiate analgesic use	May exacerbate bowel dysfunction.	High
Amiodarone (Cordarone)	Associated with QT interval problems and risk of provoking torsades de pointes. Lack of efficacy in older adults.	High
Orphenadrine (Norflex)	Causes more sedation and anticholinergic adverse effects than safer alternatives.	High
Guanethidine (Ismelin)	May cause orthostatic hypotension. Safer alternatives exist.	High
Guanadrel (Hylorel)	May cause orthostatic hypotension.	High
Cycloandelate (Cyclospasmol)	Lack of efficacy.	Low
Isoxsuprine (Vasodilan)	Lack of efficacy.	Low
Nitrofurantoin (Macrochantin)	Potential for renal impairment. Safer alternatives available.	High
Doxazosin (Cardura)	Potential for hypotension, dry mouth, and urinary problems.	Low
Methyltestosterone (Android, Virilon, and Testrad)	Potential for prostatic hypertrophy and cardiac problems.	High
Thioridazine (Mellaril)	Greater potential for CNS and extrapyramidal adverse effects.	High
Mesoridazine (Serentil)	CNS and extrapyramidal adverse effects.	High
Short-acting nifedipine (Procardia and Adalat)	Potential for hypotension and constipation.	High
Clonidine (Catapres)	Potential for orthostatic hypotension and CNS adverse effects.	Low
Mineral oil	Potential for aspiration and adverse effects. Safer alternatives available.	High
Ciimetidine (Tagamet)	CNS adverse effects including confusion.	Low
Ethacrynic acid (Edecrin)	Potential for hypertension and fluid imbalances. Safer alternatives available.	Low
Desiccated thyroid	Concerns about cardiac effects. Safer alternatives available.	High
Amphetamines (excluding methylphenidate hydrochloride and anorexics)	CNS stimulant adverse effects.	High
Estrogens only (oral)	Evidence of the carcinogenic (breast and endometrial cancer) potential of these agents and lack of cardioprotective effect in older women.	Low

Abbreviations: CNS, central nervous system; COX, cyclooxygenase; GI, gastrointestinal; NSAIDs, nonsteroidal anti-inflammatory drugs; SIADH, syndrome of inappropriate antidiuretic hormone secretion. Used with permission from: Fick DM, Cooper JW, Wade WE, et al. Updating the Beers criteria for potentially inappropriate medication use in older adults *Arch Intern Med* 2003; 163:2716-2724.

**Table 3. 2002 Criteria for Potentially Inappropriate Medication Use in Older Adults: Considering Diagnoses or Conditions**

DISEASE OR CONDITION	DRUG	CONCERN	SEVERITY RATING (HIGH OR LOW)
Heart failure	Disopyramide (Norpace), and high sodium content drugs (sodium and sodium salts [alginate bicarbonate, biphosphate, citrate, phosphate, salicylate, and sulfate])	Negative inotropic effect. Potential to promote fluid retention and exacerbation of heart failure.	High
Hypertension	Phenylpropanolamine hydrochloride (removed from the market in 2001), pseudoephedrine; diet pills, and amphetamines	May produce elevation of blood pressure secondary to sympathomimetic activity.	High
Gastric or duodenal ulcers	NSAIDs and aspirin (>325 mg) (coxibs excluded)	May exacerbate existing ulcers or produce new/additional ulcers.	High
Seizures or epilepsy	Clozapine (Clozaril), chlorpromazine (Thorazine), thioridazine (Mellaril), and thiothixene (Navane)	May lower seizure thresholds.	High
Blood clotting disorders or receiving anticoagulant therapy	Aspirin, NSAIDs, dipyridamole (Persantin), ticlopidine (Ticlid), and clopidogrel (Plavix)	May prolong clotting time and elevate INR values or inhibit platelet aggregation, resulting in an increased potential for bleeding.	High
Bladder outflow obstruction	Anticholinergics antihistamines, gastrointestinal antispasmodics, muscle relaxants, oxybutynrin (Ditropan), flavoxate (Urispas), anticholinergics, antidepressants, decongestants, and tolterodine (Detrol)	May decrease urinary flow, leading to urinary retention.	High
Stress incontinence	$\alpha$ -Blockers (Doxazosin, Prazosin, and Terazosin), anticholinergics, tricyclic antidepressants (Imipramine hydrochloride, doxepin hydrochloride, and amitriptyline hydrochloride), and long-acting benzodiazepines	May produce polyuria and worsening of incontinence.	High
Arrhythmias	Tricyclic antidepressants (Imipramine hydrochloride, doxepin hydrochloride, and amitriptyline hydrochloride)	Concern due to proarrhythmic effects and ability to produce QT interval changes.	High
Insomnia	Decongestants, theophylline (Theodur), methylphenidate (Ritalin), MAOIs, and amphetamines	Concern due to CNS stimulant effects.	High
Parkinson disease	Metoclopramide (Reglan), conventional antipsychotics, and tacrine (Cognex)	Concern due to their antidopaminergic/cholinergic effects.	High
Cognitive impairment	Barbiturates, anticholinergics, antispasmodics, and muscle relaxants. CNS stimulants: dextroAmphetamine (Adderall), methylphenidate (Ritalin), methamphetamine (Desoxyn), and pemolin	Concern due to CNS-altering effects.	High
Depression	Long-term benzodiazepine use. Sympatholytic agents: methyl dopa (Aldomet), reserpine, and guanethidine (Ismelin)	May produce or exacerbate depression.	High
Anorexia and malnutrition	CNS stimulants: DextroAmphetamine (Adderall), methylphenidate (Ritalin), methamphetamine (Desoxyn), pemolin, and fluoxetine (Prozac)	Concern due to appetite-suppressing effects.	High
Syncope or falls	Short- to intermediate-acting benzodiazepine and tricyclic antidepressants (Imipramine hydrochloride, doxepin hydrochloride, and amitriptyline hydrochloride)	May produce ataxia, impaired psychomotor function, syncope, and additional falls.	High
SIADH/hyponatremia	SSRIs: fluoxetine (Prozac), citalopram (Celexa), fluvoxamine (Luvox), paroxetine (Paxil), and sertraline (Zoloft)	May exacerbate or cause SIADH.	Low
Seizure disorder	Bupropion (Wellbutrin)	May lower seizure threshold.	High
Obesity	Olanzapine (Zyprexa)	May stimulate appetite and increase weight gain.	Low
COPD	Long-acting benzodiazepines: chlordiazepoxide (Librium), chlordiaepoxide-amitriptyline (Limbital), clidinium-chlordiaepoxide (Librax), diazepam (Valium), quazepam (Doral), halazepam (Paxipam), and chlorazepate (Tranxene). $\beta$ -blockers: propranolol	CNS adverse effects. May induce respiratory depression. May exacerbate or cause respiratory depression.	High
Chronic constipation	Calcium channel blockers, anticholinergics, and tricyclic antidepressant (Imipramine hydrochloride, doxepin hydrochloride, and amitriptyline hydrochloride)	May exacerbate constipation.	Low

Used with permission from: Fick DM, Cooper JW, Wade WE, et al. Updating the Beers criteria for potentially inappropriate medication use in older adults *Arch Intern Med* 2003; 163:2716-2724.